



CHARACTERISTICS OF THE CLINICAL COURSE OF ACUTE PANCREATITIS IN CHILDREN

Sh.M. Ibatova, F.Kh.Mamatkulova, N.B.Abdukadirova

Samarkand State Medical University, Samarkand, Uzbekistan

ABSTRACT

Acute pancreatitis is one of the most severe diseases in the pathology of the digestive system. The diagnosis of the disease is based on three signs: severe pain in the epigastrium or lumbar pain, pain that does not decrease after antispasmodic drugs, pain radiating to the heart area, behind the temporal bone, nausea, feeling no relief after vomiting, and tension in the upper abdomen. Clinical signs characteristic of the disease: acute, constant wedge-shaped pain in the epigastric region and dyspeptic disorders. The disease can also be confirmed using Shchyotkin-Blumberg, Kerte, Kach, Mayo-Robson symptoms.

KEY WORDS: *acute pancreatitis, pancreas, patients, clinical course, lumbar pain, diagnosis.*

INTRODUCTION

Today, acute pancreatitis is one of the most severe diseases in the pathology of the digestive system, and if acute pancreatitis is not diagnosed in time, it can lead to the transition to a tumor [1,6,7,9,10,13,17,20]. 4 levels of acute pancreatitis are distinguished: 1) a mild level of acute pancreatitis, in which inflammation, diffuse swelling is observed, and there is no necrosis and insufficiency; 2) moderate severity of acute pancreatitis, with transient organ failure (less than 48 hours) or pseudocysts, infiltrates, abscesses; 3) severe degree of acute pancreatitis, in which pancreonecrosis or peripancreonecrosis or persistent organ. 4) critical level of acute pancreatitis, with the development of infected pancreonecrosis or peripancreonecrosis and persistent organ failure. According to the revised Atlanta classification, there are a number of criteria for evaluating acute pancreatitis, a diagnosis is made if 2 out of 3 are present: a) a typical clinical appearance (triple sign) according to UTT: enlargement of the pancreas, decreased echogenicity, unclear contours, distribution of free fluid in the abdominal cavity; c) amylase and lipase increase in concentration by 3 times or more than normal [4,5,11,14,18].

The diagnosis of acute pancreatitis is based on three signs: severe pain in the epigastrium or lumbar pain, pain that does not decrease after antispasmodic drugs, pain radiating in the area of the heart, behind the sternum, nausea, lack of relief after vomiting, and tension in the

upper abdomen. Dry mouth, thirst, the patient's tongue is covered with a white coating. These symptoms are caused by the intake of fatty, fried and large amounts of food and diseases of the biliary tract [2,3,8,12,15,16,19].

THE PURPOSE OF THE STUDY

To determine the diagnostic criteria of acute pancreatitis.

Sources and verification methods. 25 children aged 9 to 15 years who were hospitalized in Samarkand city hospital with acute pancreatitis were surveyed. 10 (40%) were boys, 15 (60%) were girls, the average age of the patients was 12 years.

TEST RESULTS AND DISCUSSION

According to the questionnaire, 100% of patients experienced pain within the first 1.5-2 hours, and patients clearly indicated the location of pain. Of these, 59% of patients had severe, stabbing pain in the epigastric area, and 33% had pain on the left side. After 2-3 hours, pain was given to the back and spine, and in 6.3% to the left shoulder. 84% of patients had severe pain in the abdomen, and these patients turned to medical personnel on the 2-4th day of the disease. 62% of patients took painkillers at home ("no-shpa", "baralgin"), but the pain did not disappear completely, the pain decreased within 1.5-2 hours, and then severe pain started again. When we pay attention to the causes of acute pancreatitis, it was found that 35% of patients had an excessive intake of fatty, fried foods in their diet. 24% of patients had biliary dyskinesia, cholecystitis, and the remaining 31% were found to have



a genetic predisposition to the development of this disease. The clinical presentation of the disease varied, but often nausea, vomiting were observed, and patients did not feel relief afterwards. Symptoms of dry mouth, constipation, flatulence, profuse sweating, and weakness were observed in all patients. Diarrhea, high blood pressure, fainting, headache in patients signs are rarely observed. Most of the patients admitted to the department did not seek emergency medical care. Biliary dyskinesia and cholecystitis were also detected in the patients. When palpating the patients, the following signs were revealed: positive Shchyotkin-Blumberg symptom, 95% of patients also had a positive Kerte symptom (muscle tension in the projection of the pancreas and pain 5 cm above the navel), 58% of patients had a positive Kach symptom (transverse 8-11 thoracic vertebra) pain on palpation of the tumor), 50% of patients had a positive Mayo-Robson symptom (pain on palpation of the left costo-spinal angle) and 45% of patients had Mondor's triad (pain, vomiting, flatulence). Analyzing the anamnesis of patients diagnosed with acute pancreatitis, it was found that boys and girls have the same incidence of this disease. The main reason for the development of acute pancreatitis is a large and excessive intake of fatty, fried foods, biliary dyskinesia, cholecystitis, and genetic predisposition. The clinical course of the disease in patients was different. In children, the pain often started in the epigastric area or under the left rib. The pain is strong, wedge-shaped, spread to the entire abdomen within 2-3 hours. In some cases, the pain spread to the lower back, to the left shoulder, and was often lumbar. In most cases, dyspeptic changes were observed: nausea, flatulence, constipation, profuse sweating, dry mouth and thirst. Shchyotkin-Blumberg, Mayo-Robson, Kerte, Kach's symptoms, Mondor triad helped to diagnose "acute pancreatitis" in patients. These symptoms are used to confirm the diagnosis.

CONCLUSIONS

Thus, today, despite the high incidence of acute pancreatitis in children, diagnosis is insufficient. The Mondoran triad is an accurate universal method in the diagnosis of acute pancreatitis. The clinical picture characteristic of the early stage of development of acute pancreatitis is a sharp, constant wedge-shaped pain in the epigastric area, often the pain spreads to the left rib, to the surface of the abdomen, and is accompanied by dyspeptic disturbances: that is, nausea, lack of relief after vomiting,

flatulence, diarrhea. The disease can be confirmed using additional palpation methods (Shchyotkin-Blumberg, Kerte, Kach, Mayo-Robson symptoms).

REFERENCES

1. Zaporozhchenko B.S. *Changes in the activity of enzyme blood and the level of factor necrosis of the tumor in the treatment of experimental pancreatitis with the use of pentoxifylline* // *Klín. surgery.* — 2004. — No. Z. S. 1Z-14. 2. Yusupov Sh.A., Islamova D.S., Gaffarov U.B. *Structural pathology of the gastroduodenal zone and dyskinesia of the gallbladder.* // *Uchenye zapiski Orlovskogo gosudarstvennogo universiteta. Orel.* - 2014. - No. 7 (63). Chapter 2. - S.73-74
2. Islamova D.S., Gaffarov U.B., Ibatova Sh.M. *Otsenka character of kislotoproduktii u detey s yazvennoy boleznju dvenadtsatiperstnoy kishki i pervichnym chronicheskim gastroduodenitom s pomoshchyu vnutrijeludochnogo pH-metra.* // *Children's problems and children's surgery in the 21st century. Beshkek.* -2014. - S. 69-736.
3. Islamova D.S., Shadieva H.N., Togaev I.U., Ibatova Sh.M., Mamatkulova F.Kh. *Oslojnennoe technique of spring disease in a dvenadtsatiperstnoy boy and a child of senior school age: reasons, diagnostic criteria.* // *Problem biology and medicine.* -2017.- No. 2(94). - S. 51-55.
4. Dinarello C.A. *Biological basis for interleukin-1 in disease* // *Blood.* — 2004. — Vol. 87. - P. 147-167.
5. Ibatova Sh. M. *i dr. Effektivnost primeneniya abrikosovogo masla u detey s rakhitom* // *Voprosy nauki i obrazovaniya.* - 2019. - No. 27 (76). - S. 40-46.
6. Ibatova Sh. M., Mukhamadiev N. K., Akhmedov Sh. O. *Otsenka effektivnosti primeneniya abrikosovogo masla i aevita v kompleksnom lechenii detey, bolnyx rakhitom* // *Vestnik Khakasskogo gosudarstvennogo universiteta im. NF Katanova.* – 2015. – no. 12. - S. 41-43.
7. Ibatova Sh., Achilova F. *Osnovnye faktori riska razvitiya ozhireniya u detey i podrostkov* // *Journal hepato-gastroenterologicheskikh issledovaniy.* - 2021. - T. 2. – no. 3.1. - S. 2-4.
8. Ibatova Sh. M., Mamatkulova F. Kh., Ruzikulov N. Oh. *Otsenka pokazateley kletochnogo and humoralnogo faktorov immunity pri zbolevaniyax pochk u detey* // *Journal hepato-gastroenterologicheskikh issledovaniy.* – 2022. – no. SI-2.
9. Ibatova Sh., Pirmanova Sh. *Bronchoobstructive syndrome in a child at an early age: rasprostranennost, clinical manifestations, diagnosis and treatment* // *Journal problemy biologii i meditsiny.* – 2017. – no. 2 (94). - S. 174-177.



10. Islamova D. S., Ibatova Sh. M., Mamatkulova F.Kh. *O faktorakh riska razvitiya oslojnenogo techeniya yazvennoy bolezni dvenadtsatiperstnoy kishki u detey //Zdorove, demografiya, ekologiya finno-ugorskikh narodov. - 2020. - No. 4. – S. 46-47.*
11. Ibatova Sh.M., Mamatkulova F.Kh., Islamova D.S. *Modern approaches to antibiotic therapy for chronic lung diseases in children. International Journal of Multidisciplinary Research (IJMR) Volume: 8| Issue: 10| October 2022. P.139-142.*
12. Ibatova Sh. M., Mamatkulova F. Kh., Ruzikulov N.Y. *The Clinical Picture of Acute Obstructive Bronchitis in Children and the Rationale for Immunomodulatory Therapy. International Journal of Current Research and Review. Vol 12 Issue 17. September 2020. - P.152-155.*
13. Ibatova Sh. M., F. Kh. Mamatkulova, N. B. Abdukadirova, Yu. A. Rakhmonov, M. M. Kodirova. *Risk Factors for Development of Broncho-Obstructive Syndrome in Children. International Journal of Current Research and Review. Vol 12. Issue 23 December 2020.- P. 3-6.*
14. Ibatova Sh.M., Mamatkulova F.Kh., Rakhmonov Y.A., Shukurova D.B., Kodirova M.M. *Assessment of the Effectiveness of Treatment of Rickets in Children by Gas-Liquid Chromatography. International Journal of Current Research and Review. Vol 13, Issue 06, 20 March 2021. P.64-66.*
15. Islamova D.S., Ibatova Sh.M., Ibragimova Yu.B. *Osobennosti kislotoobrazuyushchey funktsii zhuldky u detey s yazvennoy boleznью dvenadtsatiperstnoy kishki. Nauchno-methodichesky magazine Dostizheniya nauki i obrazovaniya. No. 5 (85), 2022. -S.88-91.*
16. Shoir M. Ibatova, Feruza Kh. Mamatkulova, Gulnoz A. Kholikova, Dilrabo Kh. Mamatkulova. *Some indicators of lipid and phosphorus-calcium metabolism in children with rickets receiving conventional treatment. International Journal of Health Sciences, 6(S4), 2022. 3628–3638.*
17. Ibatova Sh.M., Mamatkulova F.Kh., Ibragimova M., Akhmedov D. *Clinic, diagnostics and indications for hospitalization for pneumonia in children. International Journal of Research and Development. Volume: 7, Issue:6, June 2022. P. 280-283*
18. Rau B., Cebulla M., Uhl W. *The clinical value of human pancreatic-specific protein, procarboxypeptidase B as an indicator of necrotizing pancreatitis// Pancreas. - 2004. - Vol. 17, N 2. - P. 134-139.*
19. Richter A., Nebe T., Wende K. *HLA-DR expression in acute pancreatitis // Eur. J. Surg. - 2005. -Vol. 165, N 10. - P. 947-951.*
20. Zhang J, Shahbaz M, Fang R. *Comparison of the BISAP scores for predicting the severity of acute pancreatitis in*

Chinese patients according to the latest Atlanta classification //J HepatobiliaryPancreat Sci. -2014. C.- 689-694.